Injury and Illness in Olympic Snow Sports

A Pooled Overview and Trend Analysis from Four Winter Olympic Games (2010–2022)

About the Data

Since the 2008 Beijing Summer Olympics, the International Olympic Committee (IOC) has operated a systematic injury and illness surveillance system at every edition of the Games. For Winter sports, four consecutive prospective cohort studies have recorded every new injury and illness receiving medical attention during the Games period: **Vancouver 2010, Sochi 2014, PyeongChang 2018, and Beijing 2022**.

This report serves two complementary purposes. First, **it pools all four Winter editions covering over 5,000 athlete** across 16 years of Olympic competition, producing a more stable and robust picture of the injury and illness burden than any single-edition analysis can provide. Second, it **examines how that burden has evolved across the four Games**, not to declare one edition 'safer' than another, but to identify meaningful trends that can inform ongoing prevention efforts. Focusing exclusively on snow sport disciplines: Alpine Skiing, Freestyle Skiing, Snowboarding, Cross-Country Skiing, Nordic Combined, And Ski Jumping, it offers the **snow sports community a practical reference**: which disciplines carry the highest risk, what kinds of injuries dominate, and whether the picture is improving or worsening, and where the opportunities for prevention may lie.

Evolution of Surveillance Methods

Major facts:

- Data collection shifted from paper forms (2010, 2014) to electronic reporting (2018, 2022), with National Olympic Committees (NOC) response rates consistently above 82% and reaching 99.7% in Sochi 2014.
- The core definition remained stable across all editions: *"any new or recurring condition receiving medical attention, regardless of time-loss consequences."*
- Severity classification evolved from a binary approach in 2010 (time loss yes/no, with some missing data) to a five-tier system from 2014 onwards (no time loss through >28 days), with progressively better completion.

- The 2020 International Olympic Committee (IOC) consensus statement (*from Bahr et al., published in The British Journal of Sports Medicine, 2020*) further standardised mechanism and onset definitions, adopted in Beijing 2022.
- New disciplines entered the Olympic programme over the period: slopestyle and ski halfpipe at Sochi 2014, snowboard big air at PyeongChang 2018, and ski big air and alpine mixed team parallel at Beijing 2022.

The most important consideration for pooling is data granularity:

- Vancouver 2010 reports injury location, type, and cause only at broad sport-group level ('Alpine & Snowboarding' vs. 'Nordic Skiing'), not by discipline.
- Sochi 2014 adds discipline-level incidence and severity, but not discipline-by-location or discipline-by-type matrices.
- PyeongChang 2018 is the first edition with comprehensive discipline-level matrices for all key variables, continued and refined in Beijing 2022.
- In practice, incidence data can be pooled across all four editions; the detailed injury profile (body region, type, mechanism) draws from 2018 and 2022 only.

Overall Burden

Across the four Winter Olympic Games, a total of **5,382 athlete entries** were registered in snow sport disciplines. These athletes sustained **670 injuries and 330 illnesses**, yielding a **pooled injury incidence of 12.4% injured athletes and a pooled illness incidence of 6.1% ill athletes**. In practical terms, roughly one in eight snow sport athletes sustains an injury during the Games period.


Pooled Injury Profile

Discipline Risk Hierarchy

Table 1 presents the key incidence data by discipline across all four editions, along with the pooled figures. A clear and consistent risk hierarchy emerges when we combine the data.

Discipline	Vancouver 2010		Sochi 2014		PyeongChang 2018		Beijing 2022		All Data Pooled	
	Athl. (n)	Inj. (%)	Athl. (n)	Inj. (%)	Athl. (n)	Inj. (%)	Athl. (n)	Inj. (%)	Athl. (n)	Inj. (%)
Alpine Skiing	308	14.9%	314	20.7%	322	18.0%	592	6.4%	**1536**	**13.5%**
Freestyle Ski Aerials	47	19.1%	43	48.8%	50	20.0%	49	12.2%	**189**	**24.3%**
Freestyle Ski Cross	68	19.1%	59	13.6%	57	24.6%	58	8.6%	**242**	**16.5%**
Freestyle Ski Big Air	-	-	-	-	-	-	57	28.1%	**57**	**28.1%**
Freestyle Ski Half Pipe	-	-	51	25.5%	51	27.5%	43	30.2%	**145**	**27.6%**
Freestyle Ski Moguls	57	1.8%	57	24.6%	60	5.0%	60	3.3%	**234**	**8.5%**
Freestyle Ski Slopestyle	-	-	52	30.8%	54	16.7%	58	22.4%	**164**	**23.2%**
Snowboard Big Air	-	-	-	-	69	11.6%	59	13.6%	**128**	**12.5%**
Snowboard Cross	57	35.1%	61	34.4%	70	25.7%	64	12.5%	**252**	**26.6%**
Snowboard Half Pipe	69	13.0%	66	18.2%	54	14.8%	47	8.5%	**236**	**14.0%**
Snowboard Slalom	59	6.8%	64	14.1%	62	4.8%	63	3.2%	**248**	**7.3%**
Snowboard Slopestyle	-	-	46	37.0%	66	21.2%	60	23.3%	**172**	**26.2%**
Cross-Country Skiing	292	3.1%	297	7.7%	311	5.5%	293	3.4%	**1193**	**4.9%**
Nordic Combined	52	1.9%	54	3.7%	55	1.8%	58	1.7%	**219**	**2.3%**
Ski Jumping	67	4.5%	96	7.3%	101	6.9%	103	2.9%	**367**	**5.4%**
TOTAL	**1076**	**10.7%**	**1260**	**18.1%**	**1382**	**13.3%**	**1664**	**8.6%**	**5382**	**12.4%**

The **highest-risk disciplines are consistently those characterised by aerial manoeuvres at high speed**. Freestyle Ski Big Air (28.1% pooled injured athletes, one edition at Beijing 2022), Freestyle Ski Halfpipe (27.6%), Snowboard Cross (26.6%), Snowboard Slopestyle (26.2%), Freestyle Ski Aerials (24.3%) , and Freestyle Ski Slopestyle (23.2%) **all see roughly one in four athletes injured during the Games**. Freestyle Ski Cross (16.5%) follow closely.



Alpine Skiing sits at 17.9% pooled injured athletes across 2010-2018, although this aggregate masks meaningful variation across sub-disciplines. The Beijing 2022 data, the first to report alpine events separately, shows Downhill at 16.5% of injured athletes, Giant Slalom at 9.4%, Slalom at 3.4%, and Super-G at 2.2%.

At the lower end of the spectrum, **Nordic disciplines show substantially lower injury rates**: Cross-Country Skiing at 4.9% injured athletes, Ski Jumping at 5.4%, and Nordic Combined at 2.3%. Snowboard Slalom (7.3%) and Freestyle Ski Moguls (8.5%) also sit in this lower-risk tier.

Injury Location

Note: Discipline-level injury location data is available only from PyeongChang 2018 and Beijing 2022, the two editions that published systematic body-region matrices per sport. The figures below pool these two editions. Vancouver 2010 and Sochi 2014 data are discussed separately.

Pooling the 2018 and 2022 discipline-level, the **knee is the single most commonly injured body region** in snow sports, representing approximately 12% of all injuries. This is driven primarily by Alpine Skiing, Freestyle Ski Cross, and Snowboard Cross. The ankle and hip follow at roughly 10% each. **The head accounts for approximately 9% of injuries**, with the highest concentrations in Half Pipe and Slopestyle disciplines (both Freestyle Ski and Snowboard). The shoulder/clavicle complex, lumbar spine/lower back, and hand/fingers round out the top locations.

Vancouver 2010 reported injury location at the sport-group level only ('Alpine & Snowboarding' vs. 'Nordic Skiing'). Even at this aggregate level, the **knee was the dominant location, accounting for 22.0% of all snow sport injuries**, with 24 of the 26 knee injuries occurring in the 'Alpine & Snowboarding' group. Head injuries represented 14.4% (all 17 in Alpine & Snowboarding), and shoulder/clavicle injuries 8.5%. The higher knee and head proportions in 2010 compared to the 2018–2022 pooled figures may reflect the narrower denominator (fewer disciplines, fewer low-severity injuries captured) rather than a change in injury patterns.

Sochi 2014 did not publish a systematic injury location matrix. However, the authors described severe injuries narratively: of the 61 injuries classified as severe (>7 days), 23 out of 25 ligament sprains/ruptures affected the knee, and 14 fractures were distributed across foot, hand, clavicle, ankle, face, pelvis, wrist, elbow, and ribs, which is consistent with the multi-region distribution seen in later editions.

Injury Type

Note: Discipline-level injury type data is available only from PyeongChang 2018 and Beijing 2022. The figures below pool these two editions.

Joint sprains and ligament tears are the most common injury type in the 2018 and 2022 Games, accounting for roughly 20-25% of all snow sport injuries. Separately, **three distinct contusion categories collectively form the largest injury cluster**: muscle contusions (43 cases), bone contusions (41 cases), and superficial contusions/bruises (14 cases). These are clinically distinct but together they represent **approximately 30% of the pooled injury burden**. Muscle and bone contusions are particularly common in Half Pipe and Cross disciplines (both Freestyle Ski and Snowboard). Fractures account for approximately 8-12% of injuries and are notably concentrated in Cross and Slopestyle events (both Freestyle Ski and Snowboard). **Concussions represent roughly 5-7%** of all snow sport injuries, with the highest rates in Half Pipe, Slopestyle, and Cross disciplines (both Freestyle Ski and Snowboard). Muscle strains make up about 6-8% of the injury burden.

Vancouver 2010 reported injury type at the sport-group level and showed a similar profile: sprains/ligament injuries (26.3%) and contusions (26.3%) dominated in equal measure, followed by fractures (10.2%), concussions (9.3%), and strains (6.8%). Concussions were proportionally higher in 2010 (9.3% vs. approximately 7% pooled in 2018–2022), with all 11 occurring in 'Alpine & Snowboarding' disciplines.

Sochi 2014 described severe injury types again narratively: 25 ligament sprains/ruptures (predominantly knee), 14 fractures, 11 concussions, 6 contusions, 2 muscle strains, 2 tendon ruptures, and 2 dislocations among the severe cases. While not a complete type distribution, the dominance of ligament injuries and fractures among severe cases is consistent with the data from later editions.

Injury Mechanism

Note: Discipline-level injury mechanism data is available only from PyeongChang 2018 and Beijing 2022. The figures below pool these two editions.

Contact with an object is the most common injury mechanism, accounting for approximately 25-30% of all injuries. This is the dominant mechanism in Half Pipe, Slopestyle, and Cross disciplines (both Freestyle Ski and Snowboard). **Non-contact acute trauma is the second most common mechanism** (approximately 15-20%), where the injury is caused without an external contact event. Overuse injuries with gradual onset account for roughly 10-16%, and they are more prevalent in Cross-Country Skiing.

Vancouver 2010 reported injury cause at the sport-group level, showing non-contact trauma as the single most common mechanism (31.4%), followed by contact with an object (15.3%) and field/play conditions (8.5%). The higher proportion of non-contact trauma in 2010 compared to 2018-2022 may partly reflect differences in the mechanism taxonomy used (fewer categories, broader definitions). Notably, 34 of the 37 non-contact trauma injuries occurred in the 'Alpine & Snowboarding' group (91.9%) , and 9 of 10 field/play condition injuries were also in the 'Alpine & Snowboarding' group (90%).

Sochi 2014 reported mechanism proportions at the Games-wide level: contact with an object was the most frequent mechanism (25%), followed by overuse with gradual onset (14%) and non-contact trauma (13%). Contact with the ground was the leading mechanism across most Freestyle Ski and Snowboard disciplines, consistent with the discipline-level data from later editions.


Injury Severity

Across the three editions with tiered severity data (2014, 2018, 2022), approximately 30-39% of all snow sport injuries resulted in estimated time loss of at least one day, and **13-17% were classified as severe** (more than 7 days of absence). The disciplines with the highest absolute counts of severe injuries are Snowboard Cross (21), Alpine Skiing (20), Freestyle Ski Cross (14), Snowboard Slopestyle (13), and Freestyle Ski Slopestyle (11).

By proportion of **injuries classified as severe**, Freestyle Ski Cross (35.0%), Snowboard Cross (31.3%), Freestyle Ski Moguls (30.0%), Freestyle Ski Slopestyle (28.9%), and Snowboard Slopestyle (28.9%) rank highest.

Narrative case descriptions in the individual published articles indicate that ligament ruptures (predominantly knee), fractures, and concussions are the injury types most commonly represented among severe cases across these disciplines, although the systematic data does not cross-tabulate type, severity and discipline.


When **pooling injury and severity data** from the 2018 and 2022 Winter Olympic Games, **Cross and Slopestyle events** (in both Freestyle Ski and Snowboard), together with **Freestyle Ski Halfpipe**, emerge as the disciplines with the **highest injury risk**, combining high injury incidence with high injury severity. Freestyle Ski Aerials and Freestyle Ski Big Air also show high injury incidence, although with more moderate injury severity. In contrast, Nordic Combined stands out for relatively high injury severity despite low injury incidence. At the other end of the spectrum, Cross-Country Skiing, Ski Jumping, and Snowboard Slalom show both low injury incidence and low injury severity. The bubble chart below provides a risk-weighted perspective by combining both probability and consequence.



Each discipline is positioned according to how often injuries occur (x-axis) and how severe they tend to be (y-axis), with bubble size reflects the number of athletes exposed. Disciplines located in the top-right quadrant can be interpreted as the highest-risk disciplines, as they combine both high injury incidence and high injury severity.

Training vs. Competition

Across all editions, **the majority of snow sport injuries (typically 50-65%) occur during training rather than competition**. This is one of the most actionable findings as it challenges the assumption that competition is the highest-risk moment. It may therefore redirect attention toward training environments, course familiarization sessions, and the accumulated exposure from multiple training runs.




The training dominance is most pronounced in the **acrobatic disciplines**. In Freestyle Ski Aerials, Ski Big Air, and Snowboard Slopestyle, **training accounts for roughly 70-80% of all injuries**. Alpine Skiing and the Nordic disciplines also show a training majority, although less extreme.

The main exception within snow sports is **Freestyle Ski Cross and Snowboard Cross, where competition injuries are roughly equal to or exceed training injuries**. In these disciplines, the multi-athlete racing format concentrates injury risk in competitive heats rather than individual training runs.

Sex-Stratified Injury Patterns

Pooling sex-stratified data across all four editions reveals that the overall injury incidence gap between women and men in snow sports is modest and not statistically consistent. However, several discipline-specific sex differences are practically meaningful.




Snowboard Cross shows the largest sex disparity: 40.2% of female athletes were injured across four editions, compared to 16.6% of males (a 2.4-fold difference). Freestyle Ski Slopestyle (32.9% female vs. 14.3% male) and Freestyle Ski Big Air (34.6% female vs. 19.4% male) show similar patterns with women at substantially higher injury risk. Freestyle Ski Cross (20.0% female vs. 13.4% male) and Freestyle Ski Moguls (11.3% female vs. 5.9% male) follow the same direction.

A few disciplines show the reverse pattern. In Freestyle Ski Halfpipe, male athletes have higher injury rates (32.1% vs. 22.4% female), and Snowboard Slopestyle shows near-parity (25.6% female vs. 25.5% male).

Taken together, these reversals suggest that **sex differences in injury risk are discipline-specific**, and that targeted and contextual investigation is needed.


Pooled Illness Profile

The illness picture in Olympic snow sports is dominated by **respiratory tract infections, which account for 48-70% of all illnesses** across editions – with the lower figure from Beijing 2022, where COVID-19 countermeasures suppressed respiratory transmission. Gastrointestinal complaints follow at 11-20%, and dermatological and cardiovascular issues both at 5-8%. Infection is the primary aetiology in 58-64% of all illness cases.




Discipline	Vancouver 2010		Sochi 2014		PyeongChang 2018		Beijing 2022		All Data Pooled	
	Athl. (n)	Ill. (%)	Athl. (n)	Ill. (%)	Athl. (n)	Ill. (%)	Athl. (n)	Ill. (%)	Athl. (n)	Ill. (%)
Alpine Skiing	308	4.2%	314	8.6%	322	9.6%	592	0.5%	**1536**	**4.8%**
Freestyle Ski Aerials	47	2.1%	43	11.6%	50	6.0%	49	10.2%	**189**	**7.4%**
Freestyle Ski Cross	68	2.9%	59	6.8%	57	3.5%	58	5.2%	**242**	**4.5%**
Freestyle Ski Big Air	-	-	-	-	-	-	57	1.8%	**57**	**1.8%**
Freestyle Ski Half Pipe	-	-	51	7.8%	51	11.8%	43	2.3%	**145**	**7.6%**
Freestyle Ski Moguls	57	0.0%	57	1.8%	60	5.0%	60	0.0%	**234**	**1.7%**
Freestyle Ski Slopestyle	-	-	52	5.8%	54	11.1%	58	0.0%	**164**	**5.5%**
Snowboard Big Air	-	-	-	-	69	8.7%	59	1.7%	**128**	**5.5%**
Snowboard Cross	57	10.5%	61	3.3%	70	7.1%	64	3.1%	**252**	**6.0%**
Snowboard Half Pipe	69	5.8%	66	3.0%	54	7.4%	47	2.1%	**236**	**4.7%**
Snowboard Slalom	59	6.8%	64	1.6%	62	12.9%	63	4.8%	**248**	**6.5%**
Snowboard Slopestyle	-	-	46	2.2%	66	6.1%	60	6.7%	**172**	**5.2%**
Cross-Country Skiing	292	6.8%	297	13.8%	311	11.3%	293	7.5%	**1193**	**9.9%**
Nordic Combined	52	7.7%	54	9.3%	55	10.9%	58	6.9%	**219**	**8.7%**
Ski Jumping	67	1.5%	96	2.1%	101	6.9%	103	1.0%	**367**	**3.0%**
TOTAL	**1076**	**5.1%**	**1260**	**7.8%**	**1382**	**9.1%**	**1664**	**3.1%**	**5382**	**6.1%**

Cross-country skiing carries the highest illness incidence in snow sports at 9.9% of ill athletes, followed by Nordic Combined (8.7%) and Alpine Skiing (7.5%).



The discipline-level sex-stratified illness data shows a mixed picture within snow sports. Some disciplines show a pronounced female excess: Snowboard Slalom (9.8% female vs. 1.6% male), Snowboard Big Air (11.9% vs. 0%), and Freestyle Ski Aerials (6.3% vs. 3.2%). Others show the opposite: Cross-Country Skiing, the discipline with the largest illness sample, shows a male excess (5.1% female vs. 7.1% male) across four Games, and Snowboard Half Pipe similarly shows higher male illness rates (1.9% female vs. 5.3% male). These discipline-level figures should be interpreted with caution, as several involve small absolute numbers (e.g., Snowboard Big Air: 7 female illnesses out of 59 athletes vs. 0 out of 69 males).



The **Beijing 2022 Games** provide a natural experiment: comprehensive COVID-19 countermeasures including masking, physical distancing, hand hygiene, and closed-loop management **reduced the overall snow sport illness incidence to 3.1 cases per 100 athletes**, roughly half the rate observed in Sochi 2014 (7.8 cases/100 athletes) and PyeongChang 2018 (9.1 cases/100 athletes).


Trends Across Four Winter Games

While the pooled data gives the most robust overall picture, examining how injury and illness rates have moved across the four editions reveals their patterns. A few important caveats apply: the Olympic programme expanded across editions (Freestyle Ski and Snowboard Slopestyle and Freestyle Ski Half Pipe were added at Sochi 2014; Snowboard Big Air at PyeongChang 2018; Freestyle Ski Big Air and Alpine Skiing Mixed Team Parallel at Beijing 2022), changes in reporting methodology may influence recorded rates, and the short Games period means that edition-to-edition fluctuations can reflect venue-specific conditions, weather, or simple statistical variability.

Overall Snow Sport Injury and Illness Rates

The overall snow sport **injury rate peaked at Sochi 2014 (20.1% of athletes injured)**, fell to 14.4% in PyeongChang 2018, and **dropped further to 9.2% in Beijing 2022**, the lowest rate recorded across the four editions. Vancouver 2010 sits at 11.9%.

The snow sport illness trajectory shows **a rise from 5.1% in 2010 to 7.8% in 2014 and 9.1% in 2018, followed by a big drop to 3.1% in 2022**. The 2022 decline is largely attributable to the comprehensive COVID-19 countermeasures in Beijing (masking, distancing, closed-loop management), which suppressed not only SARS-CoV-2 transmission but respiratory infections broadly.




Injury Trends

The Sochi peak may likely reflect the introduction of several high-risk disciplines (Freestyle Ski and Snowboard Slopestyle, Freestyle Ski Halfpipe) to the Olympic programme that edition, combined with a broader medical-attention definition compared to Vancouver 2010 as the surveillance system evolved. The subsequent decline should be interpreted with caution; it is unclear how much is attributable to prevention efforts versus changes in reporting methods, venue conditions, or the expanding denominator as more athletes enter lower-risk sub-disciplines.

Sport-Group Trajectories

Freestyle Skiing shows a clear arc: injury rates rose sharply from 13.4% in 2010 to 26.7% in 2014 (when Slopestyle and Halfpipe Skiing entered the programme), then declined to 18.8% in 2018 and 15.7% in 2022. This trajectory suggests that the initial introduction of acrobatic disciplines inflated the group rate, and that athletes, course designers, and rule-makers may have progressively adapted – although the rate remains substantially higher than non-acrobatic disciplines.




Snowboarding follows a similar downward trajectory from a 2014 peak (24.9%) to 15.9% in 2018 and 12.3% in 2022. The decline in **Snowboard Cross is particularly notable: from 35.1% in 2010 and 34.4% in 2014 to 25.7% in 2018 and 12.5% in 2022**.

Nordic disciplines (Cross-Country Skiing, Nordic Combined, Ski Jumping) show **consistently low and stable injury rates across all four editions**, ranging from 1.7% to 7.7%.

Alpine Skiing might present an interpretation challenge. The aggregate rate was relatively stable from 2010 to 2018 (14.9%, 20.7%, 18.0%) but dropped considerably to 6.4% in 2022. However, the 2022 figure reflects the first-time sub-discipline breakdown, when sub-disciplines are reported separately and athletes competing in multiple events are counted per event, therefore the denominator expanding and, in turn, lowering the rate. Direct comparison should be made with caution of the 2022 Alpine Skiing numbers with earlier editions.

Illness Trends

Cross-Country Skiing carries the highest illness burden throughout, peaking at 13.8% in 2014 and remaining above 7% in every edition. Nordic Combined shows a similar pattern (7.7%-10.9%).



Within Freestyle Skiing and Snowboarding, illness rates are lower and more volatile, likely reflecting the shorter competition and training exposure windows compared to endurance events.

The Beijing 2022 illness experience demonstrates that behavioural and environmental countermeasures can meaningfully reduce the illness burden.

Discipline-Level Patterns

Some individual disciplines show notable trajectories. **Freestyle Ski Half Pipe is the only discipline with a consistently rising injury rate** across all three editions since its Olympic introduction: **25.5% in 2014, 27.5% in 2018, and 30.2% in 2022**.

Freestyle Ski Aerials, as noted earlier, show extreme volatility (19.1% in 2010, 48.8% in 2014, 20.0% in 2018, 12.2% in 2022), whereas Snowboard Slopestyle, by contrast, has been more stable since its 2014 debut (37.0% in 2014, 21.2% in 2018, and 23.3% in 2022).

Key Takeaways for the Snow Sports Community

Across 5,382 athlete-entries and four Winter Olympic Games, **12.4% of snow sport athletes sustained a new injury and 6.1% an illness** during the Games period. The highest-risk disciplines are those involving aerial manoeuvres at speed: Half Pipe, Slopestyle, Big Air, and Cross events (both Freestyle Ski and Snowboard) consistently see 20-35% of athletes injured.

The **knee** is the single most commonly injured body region, and **ligament sprains/tears** are the most common injury type. The head is the second most common location, with concussions a persistent concern in Half Pipe, Slopestyle, and Cross events (both Freestyle Ski and Snowboard). **Contact with the ground or built course features** is the dominant injury mechanism.

Most injuries happen in **training**, not competition.

Snowboard Cross stands out as both a high-incidence and high-severity discipline: roughly one in four athletes is injured across four editions, with 31.3% of those injuries classified as severe.

Freestyle ski aerials present a striking volatility: the pooled incidence is 24.3%, but the edition-to-edition range swings from 12% (Beijing 2022) to 49% (Sochi 2014).

For illness, respiratory infections dominate, and the **Beijing 2022 experience** demonstrates that the practical hygiene measures employed during those Games could reasonably reduce this burden.

The **sex gap in illness incidence** turns out to be **discipline-specific**: Cross-Country Skiing and Snowboard Half Pipe show male-excess illness, while Snowboard Slalom and Snowboard Big Air show pronounced female-excess patterns, although several of these discipline-level comparisons involve small samples.

Sex-stratified injury data reveal that Snowboard Cross (40.2% female vs. 16.6% male) and Ski Slopestyle (32.9% female vs. 14.3% male) present the most extreme sex disparities.

The IOC surveillance system has matured considerably across these four editions, with each iteration capturing richer and more granular data. The 2018 and 2022 datasets in particular provide the discipline-level detail needed to design targeted prevention programmes.

Sources

- Engebretsen L, Steffen K, Alonso JM, et al. **Sports injuries and illnesses during the Winter Olympic Games 2010**. *Br J Sports Med 2010*;44:772–780.
- Soligard T, Steffen K, Palmer-Green D, et al. **Sports injuries and illnesses in the Sochi 2014 Olympic Winter Games**. *Br J Sports Med 2015*;49:441–447.
- Soligard T, Palmer D, Steffen K, et al. **Sports injury and illness incidence in the PyeongChang 2018 Olympic Winter Games**. *Br J Sports Med 2019*;53:1085–1092.
- Soligard T, Palmer D, Steffen K, et al. **Olympic Games during nationwide lockdown: sports injuries and illnesses at the Beijing 2022 Winter Olympics**. *Br J Sports Med 2023*; 0:1–8.
- Bahr R, Clarsen B, Derman W, et al. **IOC consensus statement: methods for recording and reporting of epidemiological data on injury and illness in sport 2020**. *Br J Sports Med 2020*;54:372–389.